

**Pre-Harvest Intervals, Grazing/Feeding and Crop Rotation
Restrictions for fungicides approved for use on soybeans for the
control of soybean rust in Minnesota as of April 4, 2005**

**(ALWAYS REVIEW INDIVIDUAL PRODUCT LABELS FOR CHANGES AND
COMPLETE DESCRIPTIONS OF THESE RESTRICTIONS)***

<u>Product</u>	<u>Grazing/Feeding Restrictions</u>	<u>Rotational Restrictions</u>	<u>PHI</u>
Bravo Weather Stik & Echo 720, 90DF	Do not use soybean hay or threshings from treated fields for animal feed or bedding.	None	42 days
Headline	Do not feed soybean forage for 14 days after last applica- tion. Do not feed soybean hay for 21 days after last applica- tion.	Anytime for barley, citrus dry bean, grass grown for seed, peanut, pecan, rye, sugarbeet, potato, wheat & triticale, berries, bulb vege- tables, cherry, cucurbit vege- tables, filberts, fruiting vege- tables, pistachios, root vege- tables. All others, 14 days.	21 days
Quadris	None	12 months for sorghum, oats, 14 days buckwheat, millet, rye, wild rice, non-grass animal feeds (alfalfa, clover), spices, sugar- cane. 36 days for leafy vege- tables (except Brassica) group; Brassica, leafy greens subgroup; vegetables, root subgroup; vege- tables, (tuberous and corm) sub- group; and vegetables, leaves of root and tuber group. Anytime for all others.	

<u>Product</u>	<u>Grazing/Feeding Restrictions</u>	<u>Rotational Restrictions</u>	<u>PHI</u>
Folicur	Do not feed soybean forage or hay to livestock, use for bedding, or allow livestock to graze in treated areas.	120 days except for peanut, grasses grown for seed.	30 days
Laredo EC & EW	Do not feed soybean forage and hay to livestock, use for bedding, or allow livestock to graze in treated areas.	Anytime for tomatoes, nectarines, peaches, plums, prunes asparagus, cucurbits, snap beans, almonds, grapes, peppermint, spearmint, apples, mayhaws, blackberry, currant, gooseberry, raspberry, strawberry, apricot, cherries. All others, 30 days.	28 days
Stratego	None	Anytime for corn, rice, pecans, peanuts, and wheat. 30 days for celery, cereals, sweet corn, pineapple, and sugarcane. All others 105 days.	21 days
Tilt PropiMax Bumper	Do not use hay, forage, or fodder from the soybean crop as any component of animal feed or bedding.	Anytime for celery, wheat, barley, rye, triticale, oats, corn, grasses grown for seed, peanuts, sugarcane, or rice. All others 105 days.	Through growth stage R5 (pod fill)
Domark 230	Do not graze or feed soybean forage or hay to livestock.	Anytime for soybeans or sugarbeets. All others 120 days.	Through growth stage R5 (pod fill)
Quilt	Do not use hay, forage, or fodder from the soybean crop as any component of animal feed or bedding.	Anytime for barley, wheat triticale, corn, grasses grown for seed. 45 days for oats, rye, and wild rice. All others 105 days.	Through growth stage R5 (pod fill)

<u>Product</u>	<u>Grazing/Feeding Restrictions</u>	<u>Rotational Restrictions</u>	<u>PHI</u>
Headline SBR (co-pack)	Do not feed soybean forage or hay to livestock.	Anytime for barley, citrus dry bean, grass grown for seed, peanut, pecan, rye, sugarbeet, potato, wheat & triticale, berries, bulb vegetables, cherry, cucurbit vegetables, filberts, fruiting vegetables, pistachios, root vegetables, brassica head & stem, brassica leafy greens, leafy vegetables, and leaves of root and tuber vegetables. All others, 120 days.	30 days

Minnesota Department of Agriculture, St. Paul, MN.

Reference to commercial products or trade names does not imply endorsement by the Minnesota Department of Agriculture.

List may not include all generic product names.